<image/> <section-header></section-header>		
INDEX e-WEB n°2 A(H5N1) Avian influenza – none	221	
<ul> <li>A(H5N1) Human influenza – Egypt</li> <li>"INSIDE" events: <ul> <li>Tick paralysis – Egypt</li> <li>"OUTSIDE" events: None: <ul> <li>Dengue - Djibouti</li> </ul> </li> </ul></li></ul>		
Location: World Event: A(H5N1) – Epizootic	<u>Comments</u>	
No epizooties reported this week		
Location: Egypt Event: A(H5N1) – Human	<u>Comments</u>	
<ul> <li>On June 7<sup>th</sup> 2012, the Egyptian Ministry of Health reported to <u>WHO</u> a new human case of A(H5N1) infection, in Kafr- El Sheikh governorate (cf. map 1).</li> <li>The case is:         <ul> <li>A 4 year-old girl;</li> </ul> </li> </ul>	Map 1. Kafr-El heikh governorate, Egypt.	

REPORT OF NEW HEALTH EVENTS OCCURRING <u>INSIDE</u> THE EPISOUTH AREA (Occurring in one or several EpiSouth countries)		
Location: Egypt Event: Tick paralysis	<u>Comments</u>	
<ul> <li>Mid June 2012, an article published by the <u>Journal of the Egyptian society of parasitology</u> suggested the occurrence of a tick paralysis in 4 human cases in Egypt.</li> <li>The cases, the first human cases ever reported in Egypt were 4 children living in rural area in Giza governorate: <ul> <li>The clinical pictures were confused and different clinical diagnoses were considered (rabies; myasthenia gravis; botulism; diphtheritic polyneuropathy).</li> <li>Ticks were collected (from infested children), negative clinical data and biological data were in favour of Tick paralysis.</li> <li>The encountered ticks infesting their animals were <i>Rhipicephalus sanguineus</i> on dogs, <u>_Hyalomma dromedarii_</u> on camels and <u>_Hyalomma anatolicum excavatum_</u> and <u>_Haemaphysalis_</u> sp. on goats.</li> </ul> </li> <li>Tick paralysis in human is rare and affects usually children under the age of 10.</li> <li>Tick paralysis has killed thousands of animals, mainly cows and sheep, in other parts of the world (cf. map 3); and is of concern in domestic animals and livestock in the United States.</li> </ul>	<ul> <li>Tick paralysis is the only tick-borne disease that is not caused by an infectious organism.</li> <li>The illness is caused by a neurotoxin produced in the tick's salivary gland. After prolonged attachment, the engorged tick transmits the toxin to its host. Tick paralysis results from inoculation of a toxin from tick salivary glands during a blood meal.</li> <li>The incidence of tick paralysis is unknown.</li> <li>The toxin causes symptoms within 2–7 days, beginning with weakness in both legs that progress to paralysis. The paralysis ascends to the trunk, arms, and head within hours and may lead to respiratory failure and death. If the tick is not removed, the toxin can be fatal, with reported mortality rates of 10–12 percent, usually due to respiratory paralysis.</li> <li>No vaccine is currently available for any tickborne disease, except for Tick-borne encephalitis. Individuals should therefore take precautions when entering tick-infested areas, particularly in the spring and summer months.</li> </ul>	
The public health implication is difficult to assess at this stage.		
Map 2. Worldwide distribution of Tick Paralysis in humans, livestock and pets. source: <u>http://www.lowchensaustralia.com/pests/paralysis-tick/which-ticks-cause-paralysis.htm</u>		
D. andersoni D. accidentalis D. occidentalis D. variabilis D. marginatus D. marginatus I. ricinus R. sanguineus R. nutalli R. tricuspis I. rubicundus	HUMANS LIVESTOCK PETS crenulatus	

R. evertsi R. simus

H. truncatum

WORLDWIDE DISTRIBUTION OF TICK PARALYSIS, adapted from:

Goddard, J (1988): Tick Paralysis; Infect Med 15(1):28-31, SCP communications, 1988, in Medscape article

experimental data. Canada Dept Agri Monograph No. 9, 1973.

http://www.medscape.com 1999; original article Gregson JD: Tick paralysis: An appraisal of natural and

maculatum

Medscape ®

I. holocyclus

 $\nabla$ 

REPORT OF NEW HEALTH EVENTS OCCURRING <u>OUTSIDE</u> THE EPISOUTH AREA (Not Occurring in one or several EpiSouth countries)		
Location: Djibouti Event: Dengue	<u>Comments</u>	
<ul> <li>The Ministry of Health of Djibouti reported 111 cases of "dengue-like" syndromes in Djibouti city (cf. map 3), between January and May 2012 (no further information available).</li> <li>Since November 2011, sporadic cases of dengue are reported regularly from Djibouti especially among French and Chinese expatriates.</li> <li>The serotypes DEN-1 and DEN-3 were confirmed by the French National Reference laboratory of arboviruses.</li> </ul>	<text><list-item><list-item></list-item></list-item></text>	

## Page 3/1